

# SPACE RESEARCH COORDINATION CENTER



## IONIZATION OF IONS

BY

BRIAN K. THOMAS  
DEPARTMENT OF PHYSICS  
UNIVERSITY OF PITTSBURGH

AND

J. D. GARCIA  
DEPARTMENT OF PHYSICS  
UNIVERSITY OF ARIZONA

SRCC REPORT NO. 83

UNIVERSITY OF PITTSBURGH  
PITTSBURGH, PENNSYLVANIA

14 AUGUST 1968

GPO PRICE \$ \_\_\_\_\_

CSFTI PRICE(S) \$ \_\_\_\_\_

Hard copy (HC) 3-00

Microfiche (MF) 65

ff 653 July 65

602 FORM FACILITY

N68-32562	(THRU)
(ACCESSION NUMBER)	(CODE) 25
23	(CATEGORY)
(PAGES)	
2-2620	(NASA CR OR TAX OR AD NUMBER)



The Space Research Coordination Center, established in May, 1963, has the following functions: (1) it administers predoctoral and postdoctoral fellowships in space-related science and engineering programs; (2) it makes available, on application and after review, allocations to assist new faculty members in the Division of the Natural Sciences and the School of Engineering to initiate research programs or to permit established faculty members to do preliminary work on research ideas of a novel character; (3) in the Division of the Natural Sciences it makes an annual allocation of funds to the interdisciplinary Laboratory for Atmospheric and Space Sciences; (4) in the School of Engineering it makes a similar allocation of funds to the Department of Metallurgical and Materials Engineering and to the program in Engineering Systems Management of the Department of Industrial Engineering; and (5) in concert with the University's Knowledge Availability Systems Center, it seeks to assist in the orderly transfer of new space-generated knowledge in industrial application. The Center also issues periodic reports of space-oriented research and a comprehensive annual report.

The Center is supported by an Institutional Grant (NsG-416) from the National Aeronautics and Space Administration, strongly supplemented by grants from the A. W. Mellon Educational and Charitable Trust, the Maurice Falk Medical Fund, the Richard King Mellon Foundation and the Sarah Mellon Scaife Foundation. Much of the work described in SRCC reports is financed by other grants, made to individual faculty members.

# Ionization of Ions

Brian K. Thomas  
Department of Physics  
University of Pittsburgh  
Pittsburgh, Pennsylvania

and

J. D. Garcia  
Department of Physics  
University of Arizona  
Tucson, Arizona

This technical report embodies research sponsored by the  
National Aeronautics and Space Administration under Contract number  
NGR-39-011-035 to the University of Pittsburgh, Principal Investigator  
Edward Gerjuoy.

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

# IONIZATION OF IONS<sup>\*</sup>

Brian K. Thomas<sup>†</sup>

Physics Department, University of Pittsburgh, Pittsburgh, Pennsylvania

and

J. D. Garcia

Physics Department, University of Arizona, Tucson, Arizona

## Abstract

The classical binary encounter model for ionization by charged particle impact is modified to permit evaluation of the cross section for ionization of positive ions by electron impact. General scalable expressions are obtained and compared with available experimental data and quantum theoretical approximations. The dependence on ionic charge is discussed, and a simple physical interpretation emerges. Our results indicate that this model is as reliable as the Born approximation for this process; i.e., they agree very well with experiment for energies much larger than threshold and are everywhere within a factor of two.

---



<sup>\*</sup>Supported in part by the Advanced Research Projects Agency under contract No. DA-31-124-ARO-D-440 and by the National Aeronautics and Space Administration under contract No. NGR-39-011-035 at the University of Pittsburgh.

<sup>†</sup>NASA Predoctoral Fellow.

## I. INTRODUCTION

The lack of solutions to the three-body problem presents a distinctly larger handicap in considering the ionization of ions by charged particle impact than in ionization of neutrals, because of the effects of the residual ionic field. For the neutrals, the binary encounter approximation has been found to provide a reasonable description of the phenomena; in fact, the usual first Born approximation could be considered to be of this type because only the incident particle-atomic electron interaction contributes. Furthermore, recent work<sup>1-3</sup> has indicated the utility of even a classical binary encounter approximation for charged particle ionization of neutrals.

For ionization, the primary motivation for the use of a classical binary encounter model is that it provides a simple framework for estimates, which turn out to be quite reliable at high energy and within a factor of about 2 everywhere.<sup>1-4</sup> In addition, the model has been shown<sup>5</sup> to be related to a quantum treatment. Its practical significance is greatest for multi-electron atoms (and diatomic molecules), where even the Born approximation becomes unwieldy. We have thus deemed it appropriate to provide a modification of the model to make it applicable to the reactions



In Section II we evaluate the effects of the residual ion field on the cross section within the binary encounter framework. Section III contains a comparison of our results with the available experimental data and with

quantum treatments extant. We discuss only reaction (1); the changes required for reaction (2) are straight-forward. Our results agree very well with experiment for energies much larger than threshold and are everywhere within a factor of 2. They are as reliable as the simple Coulomb-Born approximation though not as accurate as close-coupling results. Our formulation also provides some interpretational advantages.

## II. MODEL FOR ION IONIZATION

The binary encounter approximation consists of the assumption that the significant interaction is the energy exchange between the incident charged particle, of velocity  $\vec{v}_1$ , and an atomic electron of velocity  $\vec{v}_2$ . Thus the cross section for ionization of a neutral atom is

$$\sigma_{\text{ion}} = \sum_i n_i \int_{U_i}^{E_1} \sigma_{\Delta E}^{\text{eff}}(v_1, v_{2i}) d(\Delta E) \quad (3)$$

where  $\sigma_{\Delta E}^{\text{eff}}$  is the cross section for exchange of energy  $\Delta E$ , in the laboratory frame, averaged over all orientations of  $v_{2i}$ , and  $n_i$  is the number of equivalent electrons whose energy is  $U_i$ . The result (3) is to be averaged over the speed distributions of the bound electrons.

In this section we present a model for calculating the cross sections for reaction (1), taking into account the effects of the residual field of the ion. Atomic units are used throughout.

Our model can be simply stated as follows: we consider an electron with kinetic energy  $E_1$  incident on a fixed positive ion with net charge  $Z'$ . At a distance  $\xi$  from the nucleus, the incident electron undergoes an essentially binary collision with a bound electron, of binding energy  $U$ , resulting

in an energy transfer  $\Delta E \geq U$ . At the distance  $\xi$ , the incoming electron has a kinetic energy

$$E_1' = E_1 + \frac{Z'}{\xi} \geq E_1, \quad (4)$$

so that the total cross section for the energy exchange collision is given by

$$\sigma'(E_1', E_1, U) = \left\langle \int_U^{E_1} \sigma_{\Delta E}^{\text{eff}}(v_1', v_2) d(\Delta E) \right\rangle_{\text{ave}} \quad (5)$$

where  $\langle \rangle_{\text{ave}}$  denotes an averaging over the speed distribution  $f(v_2)$  of the bound electron. In Eq. (5), the upper limit of the integral must be  $E_1$ , not  $E_1'$ , since for ionization, both electrons are to be in positive energy states after collision. The total cross section for ionization will be related to  $\sigma'$  as indicated in Fig. 1. We assume that  $\sigma'$  determines an average off-axis distance  $\rho$  from the relation  $\sigma' = \pi\rho^2$ . The parameters  $\xi$  and  $\rho$  then determine a trajectory for the incident electron in the presence of the asymptotic charge  $Z'$  prior to the binary encounter. This trajectory in turn specifies the initial impact parameter  $b$  for the incident electron. The total cross section for ionization is then  $\sigma = \pi b^2$ . Our final result is presented in the form of a correction factor to an appropriate result for neutrals.

The collision radius  $\xi$  depends on both the distance of the bound electron from the nucleus  $\vec{r}_A$  and an electron-electron separation  $\vec{\delta}$  such that an energy exchange  $\Delta E \geq U$  can occur. We use an average over relative orientations,

$$\xi = |\vec{r}_A + \vec{\delta}|_{\text{ave}} = \begin{cases} \frac{1}{3r_A} [3r_A^2 + \delta^2] & \text{if } r_A > \delta \\ \frac{1}{3\delta} [3\delta^2 + r_A^2] & \text{if } r_A < \delta \end{cases} \quad (6)$$

Classically, an average  $r_A$  can be determined from the virial theorem result  $\frac{Z' + 1}{2r_A} = U$ , where  $U$  is the binding energy, at least for hydrogenic ions. We adopt this result for all cases.

$\delta$  is related only to an energy exchange collision between the two electrons. Consider the simpler case of an isolated two-electron system in which one electron is initially at rest and the other is incident with energy  $E_{1d}$ . For this situation, the minimum laboratory scattering angle  $\theta_m$ , such that a minimum energy transfer  $\Delta E = U$  may occur is given by<sup>6</sup>

$$\sin^2 \theta_m = U/E_{1d}, \quad (7)$$

corresponding to a maximum (center of mass) impact parameter,

$$s_m = \frac{1}{E_{1d}} \cot \left( \frac{\Theta_m}{2} \right), \quad (8)$$

where  $\Theta_m = 2\theta_m$  is the center of mass scattering angle. Using the center of mass orbit equation,<sup>6</sup> together with (7) and (8), we find that the largest distance of closest approach,  $d$ , such that an energy transfer of at least  $U$  can occur is

$$d = \frac{1}{E_{1d}} [(E_{1d}/U)^{1/2} + 1] \quad (9)$$



This result was derived for one electron initially at rest; if both electrons have non-zero laboratory frame velocities,  $E_{1d}$  is the relative kinetic energy. But if we average over a spherically symmetric distribution of velocities for one electron, the resultant relative energy is the total laboratory frame energy. Thus we set  $E_{1d} = E_1 - U$  in (9) and adopt this as the value of  $\delta$ :

$$\delta \equiv \frac{1}{E_1 - U} \left[ \left( \frac{E_1}{U} - 1 \right)^{1/2} + 1 \right] \quad (10)$$

Equations (10), (6), and (4) complete our specification of  $E_1'$ :

$$E_1' = \begin{cases} E_1 + \frac{3Z'r_A}{3r_A^2 + \delta^2} & r_A > \delta \\ E_1 + \frac{3Z'\delta}{3r_A^2 + \delta^2} & r_A < \delta \end{cases}$$

with  $r_A = \frac{Z' + 1}{2U}$  and  $\delta$  as defined in (10).

We now need to find the impact parameter  $b$ , such that the incident electron intercepts the "collision sphere" at an angle defined by  $\sin \theta = \rho/\xi$ , where  $\rho = (\sigma'/\pi)^{1/2}$ , as shown in Fig. 1.<sup>7</sup> Considering the ion as fixed, the trajectory of the electron is given by<sup>6</sup>.

$$\frac{1}{r} = \frac{Z'}{2E_1 b^2} \left[ 1 + \left( 1 + \frac{4E_1^2 b^2}{(Z')^2} \right)^{1/2} \cos(\theta - \theta') \right], \quad (11)$$

where

$$\cos \theta' = \left[ 1 + \frac{4E_1^2 b^2}{(Z')^2} \right]^{-1/2}.$$

Using  $r = \xi$  and  $\theta = \sin^{-1} \rho/\xi$ , together with the requirement that if  $Z' = 0$ ,  $b = \rho$  (i.e., no correction for the neutral case), we can solve (11) for  $b$ :

$$b = \frac{1}{2} \left\{ \rho + \left[ \rho^2 + \frac{2Z'}{E_1} (\xi - [\xi^2 - \rho^2]^{1/2}) \right]^{1/2} \right\}$$

The total ionization cross section, remembering the definition of  $\rho$ , is then

$$\sigma_{\text{ion}}(E_1) \equiv \pi b^2 = \frac{1}{4} \sigma' \left\{ 1 + \left[ 1 + \frac{2Z'\pi}{E_1 \sigma'} (\xi - (\xi^2 - \sigma'/\pi)^{1/2}) \right]^{1/2} \right\}^2$$

or finally, using (4) to eliminate  $\xi$  and taking advantage of the fact<sup>2</sup> that  $\sigma'$  (hence  $\sigma$ ) is a scalable function of  $E_1/U$ ,

$$\Sigma = \frac{1}{4} \Sigma' \left\{ 1 + \left[ 1 + \frac{2Z'\pi}{\beta_1 \Sigma'} \left( \frac{Z'}{\beta_1' - \beta_1} - \left[ \frac{(Z')^2}{(\beta_1' - \beta_1)^2} - \frac{\Sigma'}{\pi} \right]^{1/2} \right) \right]^{1/2} \right\}^2 \quad (12)$$

where  $\Sigma = U^2 \sigma$ ,  $\Sigma' = U^2 \sigma'$ ,  $\beta_1 = E_1/U$ , and

$$\beta_1' = \frac{E_1'}{U} = \begin{cases} \beta_1 + \frac{\frac{3}{2} Z' (Z' + 1)}{\frac{3}{4} (Z' + 1)^2 + \Delta^2} & \frac{Z' + 1}{2} > \Delta \\ \beta_1 + \frac{3 Z' \Delta}{3\Delta^2 + (\frac{Z' + 1}{2})^2} & \frac{Z' + 1}{2} < \Delta \end{cases} \quad (13)$$

$$\Delta = U\delta = \frac{1}{\beta_1 - 1} [(\beta_1 - 1)^{1/2} + 1] .$$

Equation (12) is the desired result for the cross section for removal of an electron of binding energy  $U$  from the ion whose residual charge is  $Z'$ . The total cross section for ionization of an ion is obtained by summing over all electrons in the ion. We need still specify the function defined by Eq. (5).

A few remarks about the nature of our result are in order. The factor in curly brackets in (12) represents the effect of magnification of the cross section due to the curvature of the electron's path in the residual field. The magnification is 1 when  $Z' = 0$ , as appropriate. The other difference from the model for ionization of neutrals is in requiring an increase in the incident particle energy at which the energy exchange takes place, reflected in  $E_1'$ . Thus, the result incorporates the major features of the effect of the ion field. Both of these effects are expected to be very small for reaction (2) because of the large mass differences.

We now return to the evaluation of  $\Sigma'$  or  $\sigma'$  from Eq. (5). The required expressions for  $\sigma_{\Delta E}^{\text{eff}}(v_1', v_2)$  have been given by Gerjuoy,<sup>4</sup> among others. It already involves a spherical averaging over all orientations of  $\vec{v}_2$  with respect to  $\vec{v}_1'$ . We evaluate the integral over  $(\Delta E)$  by imposing the condition  $U \leq E_1 \leq E_1'$  and taking  $E_2$  fixed but arbitrary. We have, then, the three following possibilities<sup>4</sup> (when  $E_1 \neq E_1'$ ):

$$\sigma'(E_1', E_2, U) = \int_U^{E_1} \sigma_{iii}(v_1', v_2) d(\Delta E), \quad \text{when } U \leq E_1 \leq E_1' - E_2 \leq E_1' \quad (14a)$$

$$= \int_U^{E_1' - E_2} \sigma_{iii} d(\Delta E) + \int_{E_1' - E_2}^{E_1} \sigma_i(v_1', v_2) d(\Delta E), \quad (14b)$$

$$\text{if } U \leq E_1' - E_2 \leq E_1 \leq E_1'$$

$$= \int_U^{E_1} \sigma_i(v_1', v_2) d(\Delta E), \quad \text{if } E_1' - E_2 \leq U \leq E_1 \leq E_1', \quad (14c)$$

where

$$\int^{\Delta E} \sigma_i(v_1', v_2) d(\Delta E) = -\frac{2\pi}{3} \left(\frac{v_1'}{v_2}\right) \left(\frac{1}{E_1'}\right)^2 \left\{ \left(1 - \frac{\Delta E}{E_1'}\right)^{3/2} \left(\frac{\Delta E}{E_1'}\right)^{-2} \right\}$$

and

$$\int^{\Delta E} \sigma_{iii}(v_1', v_2) d(\Delta E) = -\frac{\pi}{E_1'} \left\{ \frac{1}{3} v_2^2 (\Delta E)^{-2} + (\Delta E)^{-1} \right\}$$

with  $E_2 = \frac{1}{2} v_2^2$ , etc. Equation (14a) does not appear if  $E_1 = E_1'$ . In that case, Eqs. (14) reduce to Stabler's<sup>8</sup> result (with the appropriate changes in notation), as expected. Inserting the results of the integrations into Eqs. (14) and introducing the scaled quantities of Eqs. (12) and (13), we have

$$\Sigma'_{ion}(\beta_1', \beta_2; \beta_1) = \frac{\pi}{\beta_1'} \left\{ \frac{2}{3} \beta_2 \left(1 - \frac{1}{\beta_1^2}\right) + \left(1 - \frac{1}{\beta_1}\right) \right\}, \text{ if } 0 \leq \beta_2 \leq \beta_1' - \beta_1 \quad (15a)$$

$$= \frac{\pi}{3\beta_1'} \left\{ 2\beta_2 + 3 - \frac{3}{(\beta_1' - \beta_2)} - \frac{2}{\beta_2^{1/2}} \frac{(\beta_1' - \beta_1)^{3/2}}{\beta_1^2} \right\}, \quad (15b)$$

$$\text{if } \beta_1' - \beta_1 \leq \beta_2 \leq \beta_1' - 1$$

$$= \frac{2\pi}{3} \frac{1}{\beta_1'} \frac{1}{\beta_2^{1/2}} \left\{ (\beta_1' - 1)^{3/2} - \frac{(\beta_1' - \beta_1)^{3/2}}{\beta_1^2} \right\}, \text{ if } \beta_1' - 1 \leq \beta_2, \quad (15c)$$

where we have expressed the inequalities in (14) as inequalities on  $\beta_2 = E_2/U$ .

For ionization  $\beta_1 \geq 1$ .

Equation (15) is required to calculate  $\Sigma'(\beta_1'; \beta_1)$ . If we adopt hydrogenic speed distributions for the bound electrons

$$f(k) = \frac{32}{\pi} \frac{k^2}{(1+k^2)^4}, \quad k^2 \equiv \beta_2 \quad (16)$$

we have

$$\Sigma'_{\text{ave}}(\beta_1'; \beta_1) = \int_0^\infty \Sigma'_{\text{ion}}(\beta_1', k; \beta_1) f(k) dk. \quad (17)$$

The integral (17), using (15) and (16), results in the following expression:

$$\begin{aligned} \Sigma'(\beta_1'; \beta_1) = & \frac{32}{3\beta_1'} \left\{ \frac{(\beta_1' - 1)^{1/2}}{2\beta_1'} \left[ \frac{5}{8} - \frac{1}{4\beta_1'} - \frac{1}{\beta_1'^2} \right] \right. \\ & + \frac{(\beta_1' - \beta_1)^{1/2}}{2\beta_1(c - \beta_1)} \left[ \frac{1}{(c - \beta_1)^2} + \frac{(2 - \beta_1)}{4\beta_1(c - \beta_1)} - \frac{(2 + 3\beta_1)}{8\beta_1} \right] \\ & + \frac{5}{16} \tan^{-1}(\beta_1' - 1)^{1/2} - \frac{(3\beta_1 + 2)}{16\beta_1^2} \tan^{-1}(\beta_1' - \beta_1)^{1/2} \\ & - \frac{3(\beta_1')^{1/2}}{c^4} \ln \frac{\beta_1'^{1/2} [(\beta_1')^{1/2} + (\beta_1' - 1)^{1/2}]}{(\beta_1')^{1/2} + (\beta_1' - \beta_1)^{1/2}} \\ & + 3\beta_1' \left\{ \frac{\beta_1' - c}{6\beta_1'c} \left[ \frac{(\beta_1' - \beta_1)^{1/2}}{(c - \beta_1)^3} - \frac{(\beta_1' - 1)^{1/2}}{(\beta_1')^3} \right] \right. \\ & + \frac{5}{24} \left( \frac{\beta_1' - c}{\beta_1'c} + \frac{1}{4c^2} \right) \left[ \frac{(\beta_1' - \beta_1)^{1/2}}{(c - \beta_1)^2} - \frac{(\beta_1' - 1)^{1/2}}{(\beta_1')^2} \right] \\ & + \left( \frac{5}{16} \frac{(\beta_1' - c)}{\beta_1'c} + \frac{3}{8c^2} + \frac{1}{2c^3} \right) \left[ \frac{(\beta_1' - \beta_1)^{1/2}}{(c - \beta_1)} - \frac{(\beta_1' - 1)^{1/2}}{\beta_1'} \right] \\ & \left. \left. - \left( \frac{5}{16} \frac{(\beta_1' - c)}{\beta_1'c} + \frac{3}{8c^2} + \frac{1}{2c^3} + \frac{1}{c^4} \right) \left( \tan^{-1}(\beta_1' - 1)^{1/2} - \tan^{-1}(\beta_1' - \beta_1)^{1/2} \right) \right\} \right\} \quad (18) \end{aligned}$$

where  $c = \beta_1' + 1$ .

We note that by setting  $\beta_1' = \beta_1$ , we can obtain the averaged ionization cross section for neutral hydrogen in the binary encounter model from (18):

$$\sigma_{\text{ion};H}(E_1) = \frac{\Sigma'(\beta_1' = \beta_1)}{U_H^2}, \quad \text{where } E_1 = U_H \beta_1.$$

The numerical calculations of Kingston<sup>9</sup> are in agreement with the results obtained from the exact expression (18) with  $\beta_1' = \beta_1$ . It should be noted that this exact result is proportional to  $1/\beta_1$  as  $E_1 \rightarrow \infty$ .

### III. RESULTS AND COMPARISONS

Since both (12) and (18) are already in scaled form, the application of these results to the ionization of any ion merely requires a sum of expressions (12) for each bound electron:

$$\sigma_{\text{ion}} = \sum_i \left\{ n_i U_i^2 \Sigma(\beta_{1i}', \beta_{1i}) \right\}, \quad (19)$$

where  $n_i$  is the number of equivalent electrons having binding energy  $U_i$ ,  $\beta_{1i} = E_1/U_i$  and  $\beta_{1i}'$  is given by (13).

One test of our model is provided by a comparison with existing quantum treatments for  $\text{He}^+$ . Figure 2 shows such a comparison with the first Coulomb-Born (CB(i)) approximation of Burgess and Rudge,<sup>10</sup> and with the second Coulomb-Born (CB(ii)) calculations of Rudge and Schwartz.<sup>11</sup> The figure shows only the values explicitly calculated, not including their extrapolation to higher energies. Also shown are the experimental values of Dolder et al.<sup>12</sup> The Coulomb-Born-exchange (CBe) calculations of Rudge and Schwartz<sup>11</sup> lie very close to the experimental values, while the close coupling approximation values of Burke and Taylor<sup>13</sup> lie close to the CB(ii)

curve; neither of these is shown. It can be seen that our model gives results consistent with the simpler CB(i) approximation, but not as close to the experimental values as the more elaborate CB(ii) and CBe approximations. We point out that all of these CB approximations require extensive numerical integration, especially at higher energies.

It should be apparent that while the magnification factor in Eq. (12) was obtained in a rather direct fashion and is relatively insensitive to the parameters used in the model, the interaction energy  $E_1'$  is considerably more model dependent. Since the model only attempts plausible approximations to the exact three-body effects, we have examined various schemes for varying  $\beta_1'$ . The solid curve in Fig. 3 is our "best" result for  $\text{He}^+$ , obtained by using  $\beta_1' = \beta_1 + 2$  in (18) and using this value of  $\Sigma'$  in (12). The explicit  $\beta_1'$  dependence of (12) is still determined by (13). The dashed curve in this figure is a semi-empirical cross-section, discussed below. We note that this result lies everywhere below the CB(ii) curve of Fig. 2, except in the region from threshold to 100 eV. We denote the result obtained by using  $\beta_1' = \beta_1 + 2$  in (18) by  $\Sigma'_N$ . Table I gives the values of  $\Sigma'_N$  for the range of values of  $\beta_1$  of any practical significance.

It is apparent that this choice of the energy dependence of  $\Sigma'$  improves agreement with experiment near threshold, and goes smoothly to the unmodified result at higher energies. (As the incident energy is increased, the effects of the residual field become less important.) The effect of using  $\Sigma'_N$  simulates in some fashion exchange effects. On this basis, we suggest that the total cross section for ionization of ions can be well approximated by (19), with the use of  $\Sigma'_N$  in (12).

The dependence on residual charge  $Z'$  is illustrated in Fig. 4. There the reduced cross-section  $Q_R = (U_H^2)^{-1} \Sigma$ , where  $U_H$  is the ionization energy of hydrogen, is plotted for various values of  $Z'$ . ( $\Sigma'_N$  has been used.) It can be seen that at high energies  $\Sigma$  becomes independent of  $Z'$ , a result which can also be obtained directly from (12) and (13). The dashed curve in Fig. 4 is the "unmagnified" result; that is, it is just the reduced cross-section  $\Sigma'_N/U_H^2$ . The broken curve is the semi-empirical reduced cross-section discussed below.

The usefulness of a classical formulation is made evident here. Quantum treatments<sup>10,11</sup> obtain the same general features displayed in Fig. 4, but their interpretation is not evident. From (12) we see that the  $Z'$ -dependence of the reduced cross-section is primarily due to the curvature of the electron in the residual field of the ion. As  $Z'$  increases, the actual curvature produced by the residual field increases, but the mean distance of the bound electron from the nucleus decreases. These two effects eventually compensate each other, so that the reduced cross-section approaches a limiting curve as  $Z' \rightarrow \infty$ . The cross-section itself, of course, has additional  $Z'$ -dependence in that it is proportional to  $1/U^2$ .

Further comparisons with experiment are presented in Figs. 5 and 6. The experimental results are from Refs. 14 through 17. In each case, the upper solid curve is the direct evaluation of (19) with  $\beta_1'$  as given by (13), and the lower curve the result of using  $\Sigma'_N$  in (12), then using (13) in (12) to determine (19). The required inner shell ionization energies were taken to be Clementi's Hartree-Fock values.<sup>18</sup> Agreement with experiment is gratifying, considering the simplicity of the model, especially for the calculations involving  $\Sigma'_N$ . Even the direct model results, however, are seen to be everywhere within a factor of about 2, and much better at high energies.



Finally, we observe that an alternative choice for the electron-electron interaction cross-section,  $\Sigma'$ , can be based upon experiment. The dashed curve in Fig. 3 is the result of using  $\Sigma'_{H,exp}(\beta_1)$  for  $\Sigma'$  in (12), where  $\Sigma'_{H,exp}(\beta_1)$  has been obtained from the fit to the experimental electron-hydrogen ionization cross-sections given in Ref. 11. Similarly, the broken curve in Fig. 4 is  $Q_R(\Sigma'_{H,exp}(\beta_1))$ . Agreement with experiment is slightly improved at high energies, as expected from the results for ionization of neutrals.<sup>9</sup>

We conclude that the binary encounter model, as modified, is as reliable for predicting ionization of ions as for neutrals. Evidently, energy exchange between the incident and bound electrons is the dominant interaction occurring in this process.

#### ACKNOWLEDGMENT

We are grateful to Professor E. Gerjuoy for several enlightening discussions.

Table I.

$\beta_1 = E_1/U$	$\frac{1}{\pi} \Sigma'_N(\beta_1' = \beta_1 + 2) \quad (\text{a.u.})$
1.25	0.1137
1.50	0.1720
1.75	0.2033
2.00	0.2202
2.50	0.2320
3.00	0.2304
5.0	0.1955
7.0	0.1623
10.0	0.1274
20.0	0.0729
30.0	0.0508

## REFERENCES

1. E. Bauer and C. D. Bartky, J. Chem. Phys. 43, 2466 (1965).
2. J. D. Garcia, E. Gerjuoy, and J. E. Welker, Phys. Rev. 165, 66 (1968).
3. J. D. Garcia, J. Chem. Phys. 47, 3679 (1967).
4. E. Gerjuoy, Phys. Rev. 148, 54 (1966).
5. J. D. Garcia, Phys. Rev. 159, 39 (1967).
6. See, for example, H. Goldstein, Classical Mechanics, Addison-Wesley, Inc., Cambridge, Mass., 1950, Chapter 3.
7. The ratio  $\rho/\xi$  is strictly less than unity, if  $Z' \geq 1$ . From Eq. (6) we see that  $\xi \geq r_A$ , so that  $U\xi \geq Ur_A = \frac{Z'+1}{2}$ . Thus  $\rho/\xi \leq \frac{U\rho}{Ur_A} = \frac{2}{Z'+1} \left( U^2 \sigma'_{ion}/\pi \right)^{1/2}$ . This last factor can be shown to be less than 1 from the equations in Ref. 4.
8. R. C. Stabler, Phys. Rev. 133, A1286 (1964).
9. A. E. Kingston, Phys. Rev. 135, A1637 (1964).
10. A. Burgess and M. R. H. Rudge, Proc. Roy. Soc. (London) A273, 372 (1963).
11. M. R. H. Rudge and S. B. Schwartz, Proc. Phys. Soc. (London) 88, 563 (1966).
12. K. T. Dolder, M. F. A. Harrison, and P. C. Thoneman, Proc. Roy. Soc. (London) A264, 367 (1961).
13. P. G. Burke and A. J. Taylor, Proc. Roy. Soc. (London) A287, 105 (1965).
14. M. F. A. Harrison, K. T. Dolder, and P. C. Thoneman, Proc. Roy. Soc. (London) A274, 546 (1963).
15. M. F. A. Harrison, K. T. Dolder, and P. C. Thoneman, Proc. Phys. Soc. (London) 82, 368 (1963).
16. W. C. Lineberger, J. W. Hooper, and E. W. McDaniel, Phys. Rev. 141, 151 (1966).
17. J. W. Hooper, W. C. Lineberger, and F. M. Bacon, Phys. Rev. 141, 165 (1966).
18. E. Clementi, IBM J. of Res. & Dev. 9, 2 (1965), with suppl. tables.

## FIGURE CAPTIONS

Fig. 1. Geometry for electron-ion collision.

Fig. 2. Ionization of helium ions by electron impact. Solid curve, present results; broken curves, Coulomb-Born approximations (Ref. 10 and 11); circles, experimental results (Ref. 12).

Fig. 3. Comparison of modified binary encounter results. Solid curve, result using  $\Sigma'_N$ ; dashed curve, result using  $\Sigma'_{H, \text{exp}}$ ; circles, experiment (Ref. 12).

Fig. 4.  $Z'$ -dependence of reduced cross sections. Solid curves, reduced cross sections for various  $Z'$ ; dashed curve, "unmagnified" reduced cross section,  $\Sigma'_N/U_H^2$ ; broken curve; reduced semi-empirical cross section,  $\Sigma_{H, \text{exp}}/U_H^2$ .

Fig. 5. Electron impact ionization cross sections. In each case, the upper solid curve is the direct model result, the lower curve is the values using  $\Sigma'_N$ . (a) Neon ions; circles, experimental results (Ref. 14). (b) Nitrogen ions; circles, experimental results (Ref. 15).

Fig. 6. Electron impact ionization of alkali ions. In each case the upper solid curve is the direct model result, the lower curve is the values using  $\Sigma'_N$ . (a) Lithium ions; circles, experiment (Ref. 16). (b) Sodium ions; circles, experiment (Ref. 16). (c) Potassium ions; circles, experiment (Ref. 17).




Figure 1




Figure 2




Figure 3




Figure 4






Figure 5




Figure 6